

Hamiltonian Theory of the FQHE: Conserving Approximation for Incompressible Fractions

Ganpathy Murthy

Department of Physics and Astronomy, University of Kentucky, Lexington KY 40506

(February 6, 2008)

A microscopic Hamiltonian theory of the FQHE developed by Shankar and the present author based on the fermionic Chern-Simons approach has recently been quite successful in calculating gaps and finite temperature properties in Fractional Quantum Hall states. Initially proposed as a small- q theory, it was subsequently extended by Shankar to form an algebraically consistent theory for all q in the lowest Landau level. Such a theory is amenable to a conserving approximation in which the constraints have vanishing correlators and decouple from physical response functions. Properties of the incompressible fractions are explored in this conserving approximation, including the magnetoexciton dispersions and the evolution of the small- q structure factor as $\nu \rightarrow \frac{1}{2}$. Finally, a formalism capable of dealing with a nonuniform ground state charge density is developed and used to show how the correct fractional value of the quasiparticle charge emerges from the theory.

73.50.Jt, 05.30.-d, 74.20.-z

I. INTRODUCTION

The Fractional Quantum Hall (FQH) effect¹ has introduced us to new states of electrons in high magnetic fields². Laughlin³ explained the essence of the effect for fractions $\nu = 1/(2p+1)$ and showed that the electrons form strongly correlated states which are incompressible and have quasiparticle excitations with fractional charge³ $e^* = e/(2p+1)$.

A unified understanding of all fractions $\nu = p/(2sp+1)$ was achieved by the Composite Fermion picture of Jain⁴. In this picture, the true quasiparticles are electrons dressed by $2s$ units of statistical flux, which are called Composite Fermions (CFs). At a mean field level, the CFs then see a reduced field $B^* = B/(2sp+1)$, in which they fill p CF-Landau levels (CF-LLs), and exhibit the integer QHE. This picture has been very successful in describing the FQH states⁵.

Numerical diagonalizations of small systems^{6,7} or the use of trial wavefunctions^{5,8} are the best approaches for calculating many properties of FQH systems such as ground state energies, gaps, magnetoexciton dispersions, ground state spin transitions, etc. However, there are other interesting properties which are not accessible to these approaches, the most important of which are dynamical response functions. Partly to address these issues, and also to gain a better understanding of FQH states, several field-theoretic approaches were developed during the past fifteen years. Most of these approaches are based on the Chern-Simons (CS) transformation⁹, which performs flux attachment via the CS gauge field to obtain either bosons¹⁰⁻¹³ or fermions¹⁴. These theories have provided us with a link between the microscopic formulation of the problem and experiment, both for incompressible and compressible states^{15,16}.

Motivated by some outstanding issues in previous work¹⁷, Shankar and the present author recently developed a hamiltonian CS theory for the FQH states^{18,19}.

Inspired by the work of Bohm and Pines²⁰ on the 3D electron gas, we enlarged the Hilbert space to introduce n high-energy magnetoplasmons degrees of freedom, (n also being the number of electrons) at the same time imposing an equal number of constraints on physical states. Upon ignoring the coupling between the oscillators and the fermions we obtained some well known wavefunctions^{3,4}. A final canonical transformation was employed to decouple the fermions from the oscillators in the infrared limit.

After the final canonical transformation, we obtained the following expressions¹⁹ for the electronic charge density ρ_e and the constraint χ for $\nu = p/2p+1$ at small q :

$$\rho_e(\vec{q}) = \frac{q}{\sqrt{8\pi}} c(A(\vec{q}) + A^\dagger(-\vec{q})) + \sum_j e^{-i\vec{q}\cdot\vec{r}_j} - \frac{il^2}{1+c} \sum_j \vec{q} \times \vec{\Pi}_j e^{-i\vec{q}\cdot\vec{r}_j} \quad (1)$$

$$\chi(\vec{q}) = \sum_j e^{-i\vec{q}\cdot\vec{r}_j} + \frac{il^2}{c(1+c)} \sum_j \vec{q} \times \vec{\Pi}_j e^{-i\vec{q}\cdot\vec{r}_j} \quad (2)$$

where $A, A^\dagger$ refer to the annihilation and creation operators of the magnetoplasmon oscillators, $l = 1/\sqrt{eB}$ is the magnetic length, $\vec{\Pi}_j = \vec{P}_j + e\vec{A}^*(r_j)$ is the velocity operator of the CFs, and $c = \sqrt{2\nu}$. Instead of using the expression of eq(1) for the electronic charge density we added a multiple of the constraint to it in subsequent calculations. The rationale for doing so is that in an exact calculation in the physical subspace (which is annihilated by the constraints) using the operator $\rho_e - \xi\chi$ with *arbitrary* ξ instead of ρ_e should not change the physical answer. However, in an approximate calculation in the full space different choices of ξ will produce different answers. The choice $\xi = c^2 = 2\nu$ is particularly advantageous, as will be seen below. With this “preferred” choice, the expression for the density becomes

$$\rho_p(\vec{q}) = \frac{q}{\sqrt{8\pi}} c(A(\vec{q}) + A^\dagger(-\vec{q}))$$

$$+ \frac{1}{2p+1} \sum_j e^{-i\vec{q}\cdot\vec{r}_j} - il^2 \sum_j \vec{q} \times \vec{\Pi}_j e^{-i\vec{q}\cdot\vec{r}_j} \quad (3)$$

The oscillator piece saturates Kohn's theorem²¹. The rest, to be called $\bar{\rho}_p$, has the following properties:

- The presence of the velocity operator $\vec{\Pi}$ in $\bar{\rho}_p$ shows that the CFs respond to the effective field B^* .
- $\bar{\rho}_p$ satisfies the magnetic translation algebra (MTA)²² to lowest leading order. Since this is the algebra of the electron density in the lowest Landau level (LLL), we have performed the LLL projection correctly in the infrared.
- Note that $\bar{\rho}_p$ is a sum of a monopole with charge $e^* = e/(2p+1)$, which is the charge associated with the CF, and a dipole piece²³ which alone survives at $\nu = 1/2$. (A number of recent constructions have emphasized this dipolar aspect^{24–27}).
- We also find that as $\vec{q} \rightarrow 0$ all transition matrix elements of $\bar{\rho}_p$ from the HF ground state vanish at least as q^2 (a consequence of Kohn's theorem²¹ for incompressible states²²).

These are all nonperturbatively correct properties of the true electronic density operator in the LLL. In our formulation the full Hilbert space of the CFs is too large, and the physical LLL will be reached only after the constraints have been properly imposed. Unfortunately, we do not know how to select the physical states operationally in the final representation. The use of $\bar{\rho}_p$ represents a shortcut that obviates the need to impose constraints. The idea is that since $\bar{\rho}_p$ displays the same properties in the full space that the true electronic density is known to possess in the physical space, calculations carried out with $\bar{\rho}_p$ in the full space will hopefully provide a reasonable approximation to the true physical answers.

The Hamiltonian of the low-energy sector now is

$$H = \frac{1}{2} \int \frac{d^2q}{(2\pi)^2} v(q) \bar{\rho}_p(-q) \bar{\rho}_p(q). \quad (4)$$

where $v(q)$ is the electron-electron interaction.

Once the constraints have been used to construct $\bar{\rho}_p$, they are ignored in subsequent calculations. One assumes that the Hartree-Fock (HF) approximation is sufficient to capture the correct physics. This assumption is borne out by calculations of gaps^{28,29} and the temperature dependence of the spin polarization^{30,31}. It even provides a reasonable qualitative description of the magnetoexciton dispersions³².

Despite these successes, the above formulation has a few disadvantages. Firstly, the expressions for the density and constraint operators are known only to first order in q . The use of the small- q preferred density operator for all q renders the gap infinite for the Coulomb interaction, and one has to use a potential that is exponentially suppressed at large q in order to get finite results^{33,34}. In computing magnetoexciton dispersions

using the small- q approach³² the author used a cutoff in the number of CF-Landau levels (CF-LLs) corresponding to keeping the correct number of single-particle states in the Hilbert space to obtain finite results for the magnetoexciton dispersions for any interaction. However, for realistic potentials the dispersions were still strongly dependent on the CF-LL cutoff³², casting doubt on the robustness of the approach. A related disadvantage is that the density operator satisfies the MTA only to leading order in q .

As stated above, the preferred density operator was obtained by adding a multiple of the constraint¹⁹ to the canonical transform of the original electron density. This made certain nonperturbative features of the charge density explicit. However, it would be desirable and more satisfying to see these nonperturbative features emerging from the formalism rather than having to put them in.

Shankar³⁵ extended the previous small- q approach to all orders in q by exponentiating the first two terms of the small- q expressions for the density (eq(1)) and the constraint (eq(2)). The resulting expressions (considered in more detail in the next section) have some very desirable algebraic properties; the density operator obeys the magnetic translation algebra exactly at all q , and the density and the constraint commute. This implies that the Hamiltonian constructed from the density operator commutes with the constraint, which makes a standard conserving approximation³⁶ possible. In such an approximation, the correlator of the constraint with either the physical density or itself is explicitly zero. It is worth noting that the algebra found by Shankar reduces to that found by Pasquier and Haldane²⁵ for bosons at $\nu = 1$ (in this case $c^2 = \nu$ in the above formulas, and bosons at $\nu = 1$ are mapped to fermions in zero average field, just as electrons are at $\nu = \frac{1}{2}$). This case was treated in great detail by Read²⁶, who computed bosonic response functions in a conserving approximation.

In this paper, we will follow this conserving route to calculating response functions for the incompressible fractions. This corresponds to using the HF approximation on the Hamiltonian to calculate single-particle energies and calculating bosonic correlators in the time-dependent HF (TDHF) approximation. Several results of interest are obtained. Firstly, it is known that the spectral function of the correlator of the projected electron density operator cannot contain terms of order q^2 , since this would violate Kohn's Theorem²¹. Naively, a density perturbation produces excitons from filled states to empty states. The electronic density operator has matrix elements of order q to some of these states, and it is interesting to see how these disappear in the final correlator. It turns out that there is always one linear combination of naive excitons with a zero eigenvalue in the TDHF approximation. This corresponds in a very close sense (to be made precise later) to the constraint, and represents the unphysical sector of the theory. The electronic density operator does not couple to this unphysical sector in the TDHF approximation, and does indeed have matrix elements of order q^2 or higher to the physical excitonic states, in compliance with Kohn's the-

orem. As a consequence of the q^2 matrix elements the structure factor $S(q)$, which is the equal time correlator $\langle \rho_e(q, t) \rho_e(-q, t) \rangle$, should go as q^4 in the incompressible states. However, Read's work²⁶ shows that at $\nu = \frac{1}{2}$ this behaves as $q^3 \log q$. We will show that for the fraction $\nu = p/(2p+1)$, the coefficient of the q^4 term in the structure factor diverges (for $p > 1$) as

$$\frac{1}{2} \frac{p^4 - 3p^3 + \frac{5}{4}p^2 + 3p + \frac{7}{4}}{p^2 - 1} \quad (5)$$

This result shows that as $p \rightarrow \infty$, a power series expansion of $S(q)$ must fail, and is consistent with the above nonanalytic result²⁶.

We will also present results for the magnetoexciton dispersions for $1/3$ and $2/5$. The positions of the minima are found to be exactly at the values of ql_0 found numerically⁵⁻⁸. This attests to the fact that although the all- q formalism was derived by extending a small- q approach, it nevertheless manages to capture the structure of the magnetoexcitons up to $ql_0 \approx 1.5$.

The fractional quasiparticle charge^{3,37,38} is one of the most striking and robust features of the FQHE, and it must emerge from any theory that correctly describes the FQHE. To see how this appears in the all- q Hamiltonian theory, a further extension of the formalism is necessary. Shankar's all- q formalism³⁵ is designed to work best in a uniform background. In an inhomogeneous background there will be a nonuniform average of the electronic charge density $n(\vec{r}) = \bar{n} + \delta n(\vec{r})$, which we expect by the principles of flux attachment to lead to a nonuniform effective magnetic field. Clearly, adding a localized quasiparticle produces just such a nonuniform charge density and effective field. A version of the all- q theory applicable to first order in $\delta n/\bar{n}$ will be presented, and used to show the emergence of the fractional quasiparticle charge. Such a formalism is also useful if one wants to treat disorder or edges in a conserving approximation.

The structure of this paper is as follows: In Section II we provide a brief introduction to the Hamiltonian theory of the FQHE, including the single-particle states. In Section III we will present the structure of the conserving approximation and the details of how it is carried out. In Section IV we will analyse the TDHF matrix, show that it has zero modes corresponding to the unphysical modes. The result for the small- q structure factor for $\nu = p/(2p+1)$ will be derived in Section V. In Section VI we will present results for the magnetoexciton dispersions for $\frac{1}{3}$ and $\frac{2}{5}$ for q not too small. In Section VII we will present the extension of the all- q formalism to the case of nonuniform density, and see how the fractional quasiparticle charge emerges naturally in the theory. We end with some conclusions.

II. HAMILTONIAN THEORY OF THE FQHE

As described in the introduction, the small- q version of the Hamiltonian theory^{18,19} of the FQHE was derived

by introducing extra degrees of freedom which assumed the role of the magnetoplasmons, and supplementing the Hamiltonian with constraints. After decoupling this led to a theory with many desirable properties, but also some deficiencies, notably involving the consistency of the theory for all q .

We will take the all- q extension of this formalism as described by Shankar³⁵ as our starting point. In this formalism two sets of guiding center coordinates are associated with each CF.

$$\vec{R}_e = \vec{r} - \frac{l^2}{1+c} \hat{z} \times \vec{\Pi} \quad (6)$$

$$\vec{R}_v = \vec{r} + \frac{l^2}{c(1+c)} \hat{z} \times \vec{\Pi} \quad (7)$$

Here $c^2 = 2\nu$, and $\vec{r}$ and $\vec{\Pi}$ are the position and velocity operators of the CF. The algebra of these objects is of fundamental importance.

$$[R_{e,\alpha}, R_{e,\beta}] = -il^2 \epsilon_{\alpha\beta} \quad (8)$$

$$[R_{e,\alpha}, R_{v,\beta}] = 0 \quad (9)$$

$$[R_{v,\alpha}, R_{v,\beta}] = i \frac{l^2}{c^2} \epsilon_{\alpha\beta} \quad (10)$$

The first set of coordinates $\vec{R}_e$ is identified with the guiding center coordinates of the electron, and the second set $\vec{R}_v$ corresponds to the guiding center coordinates of an object with a magnetic algebra charge of $-e/c^2$. We will call this object the *pseudovortex* (for brevity, Shankar simply calls it the vortex^{35,31}). Upon projection to the physical subspace we expect that this coordinate will disappear, leaving behind the proper correlations between electrons, that is, vortices. Magnetic translation operators T_e , T_v and densities ρ_e , ρ_v can be formed from the two sets

$$\rho_e(\vec{q}) = e^{-q^2 l^2/4} T_e(\vec{q}) = e^{-q^2 l^2/4} \sum_i e^{-i\vec{q} \cdot \vec{R}_{e,i}} \quad (11)$$

$$\rho_v(\vec{q}) = e^{-q^2 l^2/4c^2} T_v(\vec{q}) = e^{-q^2 l^2/4c^2} \sum_i e^{-i\vec{q} \cdot \vec{R}_{v,i}} \quad (12)$$

The operator T_e obeys the algebra of the electronic magnetic translation operators projected to the LLL²²

$$[T_e(\vec{q}_1), T_e(\vec{q}_2)] = 2i \sin\left(\frac{l^2}{2} \vec{q}_1 \times \vec{q}_2\right) T_e(\vec{q}_1 + \vec{q}_2) \quad (13)$$

while the operator T_v satisfies a similar algebra

$$[T_v(\vec{q}_1), T_v(\vec{q}_2)] = -2i \sin\left(\frac{l^2}{2c^2} \vec{q}_1 \times \vec{q}_2\right) T_v(\vec{q}_1 + \vec{q}_2) \quad (14)$$

The density formed from the electronic guiding center is identified with the electronic density operator, while that formed from the pseudovortex guiding center is the constraint³⁵. It is easy to verify that the small- q limit of these expressions indeed reduce to those derived in the small- q theory (eqs(1,2)). Since they commute with each other, the Hamiltonian formed from ρ_e alone will commute with the constraint.

In previous uses of this all- q formalism³¹, the *preferred* density $\rho_p = \rho_e - c^2 \rho_v$ was used. The preferred density has many desirable features, as described in the introduction, and gives numbers in good agreement with

numerical and experimental results for many physical properties^{28,30,31}. However, in addressing issues of principle, such as the form of the density correlator at small- q for $\nu = 1/2$ ^{39,27}, naive calculations based on the preferred density fail to produce the correct answer, and a conserving calculation^{36,26} is needed.

Before we can compute density correlators, we need to construct the single-particle states of the theory. Our Hamiltonian is

$$H = \int \frac{d^2q}{(2\pi)^2} v(q) \rho_e(\vec{q}) \rho_e(-\vec{q}) \quad (15)$$

From eq(6) we see that the coordinate R_e contains $\vec{\Pi}$, the velocity operator in the effective field. Thus, Landau level states in the effective field make a natural basis in which to represent the various operators. In the Landau gauge, these states are

$$\phi_{n,X} = C e^{iXy/(l^*)^2} e^{-(x-X)^2/2(l^*)^2} H_n((x-X)/l^*) \quad (16)$$

where C is a normalization constant, H_n are the Hermite polynomials, $l^* = 1/\sqrt{eB^*}$ is the magnetic length in the effective field, n represents the Composite Fermion-Landau Level (CF-LL) index, and $X = 2\pi j(l^*)^2/L$ is the degeneracy index (the degeneracy of each CF-LL is $\frac{L^2}{2\pi(l^*)^2}$). In terms of these states the translation operators can be represented as follows:

$$T_e(\vec{q}) = \sum_{\{n_i\}X} e^{-iq_x X} d_{n_1,X - \frac{Qy l^*}{2}}^\dagger d_{n_2,X + \frac{Qy l^*}{2}} \rho_{n_1 n_2}(\vec{q}) \quad (17)$$

$$T_v(\vec{q}) = \sum_{\{n_i\}X} e^{-iq_x X} d_{n_1,X - \frac{Qy l^*}{2}}^\dagger d_{n_2,X + \frac{Qy l^*}{2}} \chi_{n_1 n_2}(\vec{q}) \quad (18)$$

where the matrix elements are

$$\rho_{n_1 n_2}(\vec{q}) = (-1)^{n_< + n_2} \sqrt{\frac{n_<!}{n_>!}} e^{i(n_1 - n_2)(\theta_q - \pi/2)} \\ (cQ/\sqrt{2})^{|n_1 - n_2|} e^{-c^2 Q^2/4} L_{n_<}^{|n_1 - n_2|} (c^2 Q^2/2) \quad (19)$$

$$\chi_{\nu}(\vec{q}) = (-1)^{n_< + n_2} \sqrt{\frac{n_<!}{n_>!}} e^{i(n_1 - n_2)(\theta_q - \pi/2)} \\ (Q/c\sqrt{2})^{|n_1 - n_2|} e^{-Q^2/4c^2} L_{n_<}^{|n_1 - n_2|} (Q^2/2c^2) \quad (20)$$

Here $c = \sqrt{2\nu}$, $Q = ql^*$, $n_<$ ($n_>$) is the lesser (greater) of n_1 , n_2 , and L_n^k are the Laguerre polynomials.

It can be shown that the CF-LL states are in fact Hartree-Fock (HF) states of the Hamiltonian^{32,31}, that is, they are the best possible Slater determinantal states for this problem. The single-particle HF energies are

$$\epsilon(n) = \frac{1}{2} \int \frac{d^2q}{(2\pi)^2} v(q) e^{-q^2 l^2/2} \sum_m (1 - 2N_F(m)) |\rho_{nm}(\vec{q})|^2 \quad (21)$$

Note that the energy depends only on the CF-LL index, and not on the degeneracy index. The first term corresponds to the Hartree energy, while the second term is the Fock energy. When the preferred density is used the Hartree contribution is nontrivial, and depends on n . However, when the electron density of eq(11) is used, as we do here, the Hartree energy is a constant independent of n (the methods of the Appendix can be adapted to show this).

III. THE CONSERVING APPROXIMATION

To illustrate the idea of a conserving approximation³⁶, let us go back to the Hamiltonian constructed from the all- q density operator $\rho_e(\vec{q})$, eq(15). This should be supplemented by constraints that demand that the pseudovortex density should annihilate all physical states

$$\rho_v(\vec{q}) |\Psi_{phys}\rangle = 0 \quad (22)$$

Now define a time-ordered Green's function $G_{ve}(q, t)$ as follows (one can similarly define G_{ee} and G_{vv})

$$G_{ve}(\vec{q}, t - t') = \begin{aligned} & -i < T \rho_v(\vec{q}, t) \rho_e(-\vec{q}, t') > \\ & = -i \Theta(t - t') < \rho_v(\vec{q}, t) \rho_e(-\vec{q}, t') > \\ & \quad - i \Theta(t' - t) < \rho_e(-\vec{q}, t') \rho_v(\vec{q}, t) > \end{aligned} \quad (23)$$

where T is the time ordering operator. Consider the time development of this Green's function

$$-i \frac{\partial}{\partial t} G_{ve}(\vec{q}, t - t') = -\delta(t - t') < [\rho_v(\vec{q}, t), \rho_e(-\vec{q}, t')] > \\ -i < T[H, \rho_v(\vec{q}, t)] \rho_e(-\vec{q}, t') > \quad (24)$$

Since ρ_v and ρ_e commute, ρ_v also commutes with H . One immediately sees that G_{ve} is a constant. If one sets the constraint to zero initially, then it remains zero, and all its correlators also remain zero.

The above is true as an exact statement. However, in practice one uses some approximation scheme to calculate the Green's function. In our case, an approximation that respects $G_{ve} = G_{vv} = 0$ will be conserving. Let us see what a natural approximation scheme might be. Consider the time-ordered Green's function $G_{ee}(\vec{q}, t)$ defined as

$$G_{ee}(\vec{q}, t - t') = -i < T \rho_e(\vec{q}, t) \rho_e(-\vec{q}, t') > \quad (25)$$

and look at its time development. We have

$$i \frac{\partial}{\partial t} G_{ee}(\vec{q}, t) = \delta(t) < [\rho_e(\vec{q}, t), \rho_e(-\vec{q}, 0)] > \\ -i < T[H, \rho_e(\vec{q}, t)] \rho_e(-\vec{q}, 0) > \quad (26)$$

From the commutation relations of ρ_e with itself (eq(13)), it can be seen that a Green's function involving three densities will arise. Further time derivatives will involve higher-order density correlators. The natural way to truncate this hierarchy is to make a mean-field approximation at some stage that reduces a product of densities to a single density. One of the simplest of such approximations³⁶ reduces $[H, \rho_e]$, which is a product of four fermi operators, to a product of only two, by using the averages

$$d_{\alpha_1}^\dagger d_{\alpha_2}^\dagger d_{\beta_2} d_{\beta_1} \rightarrow < d_{\alpha_1}^\dagger d_{\beta_1} > d_{\alpha_2}^\dagger d_{\beta_2} + < d_{\alpha_2}^\dagger d_{\beta_2} > d_{\alpha_1}^\dagger d_{\beta_1} \\ - < d_{\alpha_1}^\dagger d_{\beta_2} > d_{\alpha_2}^\dagger d_{\beta_1} - < d_{\alpha_2}^\dagger d_{\beta_1} > d_{\alpha_1}^\dagger d_{\beta_2} \quad (27)$$

Here $< d_{\alpha}^\dagger d_{\beta} > = \delta_{\alpha\beta} N_F(\alpha)$, where $N_F(\alpha)$ is the Fermi occupation of the single-particle state α .

Using the HF states and their occupations in the above truncation is known as the time-dependent HF (TDHF) approximation. In the electron gas it is known that this approximation is conserving (in that case the conservation laws to be satisfied result from gauge invariance³⁶). Read²⁶ has shown that this approximation is conserving for the $\nu = 1$ boson problem as well. We will explicitly show below that TDHF is conserving for all principal fractions.

Let us proceed to some more explicit details.

A. TDHF Formalism

The final all- q theory has mapped the original strongly correlated electron problem into a weakly correlated CF problem with constraints. This CF problem has LL structure in the effective field, and integer filling for CFs. All the techniques that have been brought to bear on the problem of computing magnetoexcitons in the integer quantum Hall (IQH) states can be applied to this problem. The computation of magnetoexciton dispersions has a long history⁴⁰ in the IQHE. Many different approximations^{40,41} have been employed to obtain these dispersions in the IQHE, most of which can be subsumed into a unified TDHF treatment^{42,43}. In this approach one allows the hamiltonian to scatter particle-hole excitations with different Landau-level indices into each other, and diagonalizes the resulting matrix. As mentioned before, this method has been applied to the FQHE using the small- q hamiltonian approach³². Most of the details can be found there, so the following will touch on the highlights of the formalism.

The TDHF matrix can be derived from many different points of view. We will find it most convenient to follow the equation of motion approach of Rickayzen⁴⁴. We start with the magnetoexciton operator

$$O_{m_1 m_2}(\vec{q}) = \sum_X e^{-iq_x X} d_{m_1, X - \frac{Q_y l^*}{2}}^\dagger d_{m_2, X + \frac{Q_y l^*}{2}} \quad (28)$$

This represents a CF-particle-hole excitation with a definite momentum $\vec{q}$. To avoid a proliferation of indices we will use the shorthand $\mu = m_1 m_2$ (and sometimes $\nu = n_1 n_2$, not to be confused with the filling factor!). Now we define a time-ordered Green's function of two of these operators

$$G(\mu; \mu'; \vec{q}; t - t') = -i < T O_\mu(\vec{q}, t) O_{\mu'}(-\vec{q}, t') > \quad (29)$$

Taking the time derivative we obtain

$$\begin{aligned} -i \frac{\partial}{\partial t} G(\mu; \mu'; \vec{q}; t) &= -i < T [H, O_\mu(\vec{q}, t)] O_{\mu'}(-\vec{q}, 0) > \\ &\quad - \delta(t) < [O_\mu(\vec{q}, t), O_{\mu'}(-\vec{q}, 0)] > \end{aligned} \quad (30)$$

Now one Fourier transforms G to convert it to a function of frequency ω according to

$$G(t) = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} e^{i\omega t} G(\omega) \quad (31)$$

The average value of the equal time commutator in the ground state (the final term of eq(30)) can be carried out to give

$$< [O_\mu(\vec{q}), O_{\mu'}(-\vec{q})] > = \frac{L^2}{2\pi(l^*)^2} \delta_{m_1 m'_2} \delta_{m_2 m'_1} \times (N_F(m_1) - N_F(m_2)) \quad (32)$$

Now the Hamiltonian is expressed as a four-fermi operator

$$H = \frac{1}{2L^2} \sum_{\vec{q}, \{X_i\} \{n_i\}} v(q) e^{-iq_x(X_1 - X_2)} \rho_{n_1 n_2}(\vec{q}) \rho_{n_3 n_4}(-\vec{q}) d_{n_1, X_1 - \frac{Q_y l^*}{2}}^\dagger d_{n_2, X_1 + \frac{Q_y l^*}{2}} d_{n_3, X_2 + \frac{Q_y l^*}{2}} d_{n_4, X_2 - \frac{Q_y l^*}{2}} \quad (33)$$

One performs the commutator $[H, O_\mu]$, and in the TDHF approximation one contracts the resulting four-fermi operator to give a two-fermi operator. After the sums over degeneracy indices are performed one finds

$$\begin{aligned} [H, O_\mu(\vec{q})] &= (\epsilon(m_1) - \epsilon(m_2)) O_\mu(\vec{q}) + \\ & (N_F(m_2) - N_F(m_1)) \sum_\nu O_\nu(\vec{q}) \left(\frac{v(q)}{2\pi(l^*)^2} e^{-q^2 l^2 / 2} \rho_{n_1 n_2}(\vec{q}) \times \right. \\ & \left. \rho_{m_2 m_1}(-\vec{q}) - \int \frac{d^2 s}{(2\pi)^2} v(s) e^{-s^2 l^2 / 2} \rho_{n_1 m_1}(\vec{s}) \rho_{m_2 n_2}(-\vec{s}) e^{i\vec{s} \times \vec{Q}} \right) \end{aligned} \quad (34)$$

One can now define a vector $\Psi(\mu)$, which corresponds to the operator

$$O_\Psi = \sum_\mu \Psi(\mu) O_\mu \quad (35)$$

Thus the action of commuting with H on O_μ in the TDHF approximation can be represented as the right-multiplication of Ψ of a matrix $\mathcal{H}(\mu; \nu; \vec{q})$ in the space of these vectors

$$\begin{aligned} \mathcal{H}(\mu; \nu; \vec{q}) &= \delta_{m_1 n_1} \delta_{m_2 n_2} (\epsilon(m_1) - \epsilon(m_2)) + \\ & (N_F(m_2) - N_F(m_1)) \left(\frac{v(q)}{2\pi(l^*)^2} e^{-q^2 l^2 / 2} \rho_{n_1 n_2}(\vec{q}) \rho_{m_2 m_1}(-\vec{q}) \right. \\ & \left. - \int \frac{d^2 s}{(2\pi)^2} v(s) e^{-s^2 l^2 / 2} \rho_{n_1 m_1}(\vec{s}) \rho_{m_2 n_2}(-\vec{s}) e^{i\vec{s} \times \vec{Q}} \right) \end{aligned} \quad (36)$$

It is clear that finding the eigenvalues and eigenvectors of this matrix will also enable us to solve for the Green's function. Assume that one has found the right and left eigenvectors and corresponding eigenvalues, labelled by α

$$\mathcal{H}(\mu; \nu; \vec{q}) \Psi_\alpha^R(\nu; \vec{q}) = E_\alpha(q) \Psi_\alpha^R(\nu; \vec{q}) \quad (37)$$

$$\Psi_\alpha^L(\mu; \vec{q}) \mathcal{H}(\mu; \nu; \vec{q}) = E_\alpha(q) \Psi_\alpha^L(\mu; \vec{q}) \quad (38)$$

where sums over repeated indices are implicit.

We can choose to normalize the eigenvectors in the conventional way

$$\Psi_\alpha^L(\mu; \vec{q}) \Psi_\beta^R(\mu; \vec{q}) = \delta_{\alpha\beta} \quad (39)$$

If the matrix has a complete set of eigenvectors (we will see in the next subsection that this is not true, and

see how to deal with this situation) then one can expand $(\omega - \mathcal{H})^{-1}$ as

$$(\omega - \mathcal{H})^{-1}(\mu; \nu; \vec{q}) = \sum_{\alpha} \Psi_{\alpha}^R(\mu; \vec{q}) \frac{1}{\omega - E_{\alpha}} \Psi_{\alpha}^L(\nu; \vec{q}) \quad (40)$$

Now the solution to the Green's function can be expressed as

$$G(\mu; \mu'; \vec{q}; \omega) = \frac{L^2}{2\pi(i^*)^2} \sum_{\alpha} \Psi_{\alpha}^R(\mu; \vec{q}) \frac{1}{\omega - E_{\alpha}} \Psi_{\alpha}^L(m'_2 m'_1; \vec{q}) (N_F(m'_1) - N_F(m'_2)) \quad (41)$$

IV. FORMAL ANALYSIS OF THE TDHF MATRIX

In this section we will first show that there is a definite connection between the right and left eigenvectors, and that the eigenvalues of $\mathcal{H}$ always come in pairs $\pm E_{\alpha}$. Next we will show that there is always one eigenvector of $\mathcal{H}$ with a zero eigenvalue, and that this corresponds to the constraint. There is also another zero eigenvalue which does not correspond to any eigenvector, and the matrix $\mathcal{H}$ is *defective*, that is, does not have a complete set of states. We will see that we can nevertheless compute all the physical Green's functions, since they do not couple to the unphysical sector.

A. Two Useful Properties of $\mathcal{H}$

It is easy to see that $\mathcal{H}$ is neither a symmetric nor a Hermitian matrix, so in general one should expect right and left eigenvectors which are not Hermitian adjoints of each other. However, there is a definite connection between the right and left eigenvectors with the same eigenvalue:

$$\Psi_{\alpha}^R(\nu; \vec{q}) \propto (N_F(n_2) - N_F(n_1)) (\Psi_{\alpha}^L(\nu; \vec{q}))^* \quad (42)$$

This can easily be verified by noting that $\rho_{n_1 n_2}(\vec{q}) = (\rho_{n_2 n_1}(-\vec{q}))^*$, and taking the complex conjugate of the eigenvalue equation for the left eigenvector (eq(38)). In fact, any multiple of the above will form a good right eigenvector. However, to satisfy the normalization condition eq(39), we need to take

$$\Psi_{\alpha}^R(\nu; \vec{q}) = \text{sgn}(E_{\alpha}) (N_F(n_2) - N_F(n_1)) (\Psi_{\alpha}^L(\nu; \vec{q}))^* \quad (43)$$

The second property is that eigenvalues occur in equal and opposite pairs $\pm E_{\alpha}$. By focusing on the direct term, it is easy to verify that the phase of a term in $\mathcal{H}$ is

$$\frac{\mathcal{H}_d(\mu; \nu; \vec{q})}{|\mathcal{H}_d(\mu; \nu; \vec{q})|} = (-1)^{n_{<} + m_{<} + m_1 + n_1} e^{i(\theta_q - \pi/2)(n_1 + m_2 - n_2 - m_1)} (N_F(m_2) - N_F(m_1)) \quad (44)$$

It is clear from eq(36) that μ such that $N_F(m_2) = N_F(m_1)$ are not coupled to any other modes, and therefore have a time-evolution given solely by their energy $\epsilon(m_1) - \epsilon(m_2)$. The nontrivial part of $\mathcal{H}$ consists of those vectors for which $N_F(m_2) - N_F(m_1) \neq 0$. We can divide this subspace into a positive energy part with $N_F(m_2) - N_F(m_1) > 0$, and a negative energy part with $N_F(m_2) - N_F(m_1) < 0$. Clearly, for the trivial part of the spectrum, one can immediately construct a negative energy mode $O_{m_2 m_1}$ for every positive energy mode $O_{m_1 m_2}$.

Before we proceed, let us note that rotational invariance guarantees that the eigenvalues $E_{\alpha}(q)$ depend only on the magnitude of $\vec{q}$. In order to analyse the structure of $\mathcal{H}$ the choice $\theta_q = 0$ is particularly convenient. With this choice, it is easy to see that

$$\mathcal{H}_d(m_1 m_2; n_1 n_2; q_x) = -\mathcal{H}_d(m_2 m_1; n_2 n_1; q_x) \quad (45)$$

With a little more effort it is possible to show that the same relation holds for the exchange part, and thus for the entire matrix $\mathcal{H}$, which has the following structure

$$\mathcal{H} = \begin{pmatrix} \mathcal{H}_{++} & \mathcal{H}_{+-} \\ -\mathcal{H}_{+-} & -\mathcal{H}_{++} \end{pmatrix} \quad (46)$$

It follows that if

$$(u_+ \ u_-) \mathcal{H} = E(u_+ \ u_-) \quad (47)$$

is a left eigenvector of $\mathcal{H}$ with eigenvalue E , then

$$(u_- \ u_+) \mathcal{H} = -E(u_- \ u_+) \quad (48)$$

is a left eigenvector with eigenvalue $-E$. The only exception that could occur is if $E = 0$, in which case $(u_- \ u_+)$ might possibly be the same as $(u_+ \ u_-)$, and one is left with only a single eigenvector. This case is realized in our conserving approximation, and we now turn to its analysis.

B. Unphysical States in $\mathcal{H}$

The most important property of $\mathcal{H}$ is that it always has at least one left eigenvector with zero eigenvalue, namely

$$\Psi_0^L(m_1 m_2; \vec{q}) = \chi_{m_1 m_2}(\vec{q}) \quad (49)$$

The requisite formalism to show this explicitly is detailed in Appendix I, from which we borrow two identities eq(136)

$$\sum_m \chi_{m_1 m}(\vec{q}_1) \rho_{m m_2}(\vec{q}_2) = \langle m_1 | e^{-i(\frac{\vec{q}_1}{c} + c\vec{q}_2) \cdot \vec{R}_{CF}^c} | m_2 \rangle e^{-\frac{i}{2} \vec{Q}_1 \times \vec{Q}_2} \quad (50)$$

and eq(137)

$$\sum_m \chi_{m_1 m}(\vec{q}_1) \rho_{m m_2}(\vec{q}_2) = e^{-i \vec{Q}_1 \times \vec{Q}_2} \sum_m \rho_{m_1 m}(\vec{q}_2) \chi_{m m_2}(\vec{q}_1) \quad (51)$$

Let us now consider

$$\begin{aligned}
& \sum_{m_1 m_2} \chi_{m_1 m_2}(\vec{q}) \mathcal{H}(m_1 m_2; n_1 n_2; \vec{q}) = (\epsilon(n_1) - \epsilon(n_2)) \chi_{n_1 n_2}(\vec{q}) \\
& + \sum_{m_1 m_2} (N_F(m_2) - N_F(m_1)) \\
& \frac{v(q)}{2\pi(l^*)^2} e^{-q^2 l^2/2} \chi_{m_1 m_2}(\vec{q}) \rho_{m_2 m_1}(-\vec{q}) \rho_{n_1 n_2}(\vec{q}) \\
& - \sum_{m_1 m_2} (N_F(m_2) - N_F(m_1)) \int \frac{d^2 s}{(2\pi)^2} v(s) e^{-\frac{s^2 l^2}{2}} \\
& \rho_{n_1 m_1}(\vec{s}) \chi_{m_1 m_2}(\vec{q}) \rho_{m_2 n_2}(-\vec{s}) e^{i\vec{s} \times \vec{Q}} \quad (52)
\end{aligned}$$

Let us consider the direct and exchange terms separately. In the direct term, one m index can always be summed freely, while the other is constrained by the Fermi occupation factor N_F . The sum over the free m gives, according to eq(50)

$$\begin{aligned}
& \sum_{m_2} N_F(m_2) \rho_{n_1 n_2}(\vec{q}) \langle m_2 | e^{-i(\frac{1}{c}-c)\vec{q} \cdot \vec{R}_{CF}^c} | m_2 \rangle \\
& - \sum_{m_1} N_F(m_1) \rho_{n_1 n_2}(\vec{q}) \langle m_1 | e^{-i(\frac{1}{c}-c)\vec{q} \cdot \vec{R}_{CF}^c} | m_1 \rangle \quad (53)
\end{aligned}$$

The two terms are immediately seen to cancel. Now let us turn to the exchange terms, and consider the one that has the factor $N_F(m_2)$, and a free sum over m_1 . In this term, one can use eq(51) to exchange the ρ and χ matrix elements to obtain

$$\begin{aligned}
& - \sum_{m_2} N_F(m_2) \sum_{m_1} \chi_{n_1 m_1}(\vec{q}) \times \\
& \int \frac{d^2 s}{(2\pi)^2} v(s) e^{-\frac{s^2 l^2}{2}} \rho_{m_1 m_2}(\vec{s}) \rho_{m_2 n_2}(-\vec{s}) \quad (54)
\end{aligned}$$

Notice that the phase factor $e^{i\vec{s} \times \vec{Q}}$ has been cancelled by an opposite phase factor from eq(51). Now the angular $\vec{s}$ integral forces $m_1 = n_2$ for a rotationally invariant potential, and the result contains the Fock energy of the state n_2

$$\epsilon^F(n_2) \chi_{n_1 n_2}(\vec{q}) \quad (55)$$

Similarly, the other exchange term proportional to $N_F(m_1)$ ends up giving $-\epsilon^F(n_1) \chi_{n_1 n_2}(\vec{q})$. Due to the peculiar nature of the Hamiltonian the Hartree energy is a constant independent of the CF-LL index, and the difference of the Fock energies is the same as the difference of the full HF energies. Thus, the exchange contributions cancel the diagonal term $(\epsilon(n_1) - \epsilon(n_2)) \chi_{n_1 n_2}(\vec{q})$, and $\chi_{n_1 n_2}(\vec{q})$ is indeed an left eigenvector with zero eigenvalue for $\mathcal{H}$, as claimed.

If we try to obtain another eigenvector with zero eigenvalue using eqs(47,48), we are foiled by the fact that $\chi_{n_1 n_2}(q_x) = \chi_{n_2 n_1}(q_x)$, which gives us back the same eigenvector. The matrix $\mathcal{H}$ is *defective*, that is, it does not have a complete set of eigenvectors. The null subspace of $\mathcal{H}$ is two-dimensional, one of the vectors being the eigenvector $\Psi_0^L(m_1 m_2; \vec{q}) = \chi_{m_1 m_2}(\vec{q})$. The other vector (call it Ψ_0^R) is such that $\Psi_0^L \mathcal{H} = \Psi_0^L$. In other words, in this subspace, $\mathcal{H}$ looks like

$$\begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} \quad (56)$$

Fortunately, this singular behavior of the matrix is not a problem in computing physical correlators, because the density does not couple to the null subspace, as will be seen explicitly in the small q limit in the next section. To compute physical correlators, one need only project out this two-dimensional null subspace, and then the matrix inverse $(\omega - \mathcal{H})^{-1}$ looks exactly as in eq(41), except that the sum is only over physical eigenvalues.

V. SMALL Q ANALYSIS OF $\mathcal{H}$

The goal of this section is to compute the small- q behavior of the structure factor $S(q)$, which is the equal-time density-density correlator in the ground state. To this end, we will carry out an analysis of $\mathcal{H}$ for small q , and show that one needs to keep at most a 6×6 matrix to get the structure factor correct to order q^4 . One of the results of this section will be that the coefficient of this q^4 term diverges as $p \rightarrow \infty$, showing that the limits $q \rightarrow 0$ and $p \rightarrow \infty$ do not commute.

Noting that the density can be expressed as

$$\rho_e(\vec{q}) = \sum_{m_1 m_2} e^{-q^2 l^2/2} \rho_{m_1 m_2}(\vec{q}) O_{m_1 m_2}(\vec{q}) \quad (57)$$

we can write the density-density correlator $S(\vec{q}, \omega) = \langle \rho_e(\vec{q}, \omega) \rho_e(-\vec{q}, -\omega) \rangle$ as

$$\begin{aligned}
S(\vec{q}, \omega) &= \frac{L^2}{2\pi(l^*)^2} e^{-q^2 l^2/2} \\
& \sum_{\alpha, m_1 m_2} \rho_{m_1 m_2}(\vec{q}) \Psi_\alpha^R(\mu; \vec{q}) \frac{1}{\omega - E_\alpha + i\epsilon \text{sgn}(E_\alpha)} \\
& \Psi_\alpha^L(n_1 n_2; \vec{q}) (N_F(n_2) - N_F(n_1)) \rho_{n_2 n_1}(-\vec{q}) \quad (58)
\end{aligned}$$

where the sum is over physical states only, and the $i\epsilon$ factors appropriate to the time-ordered Green's function have been inserted. Now one can go back to the time-domain Green's function, and take the limit of equal time to write in a transparent notation

$$\begin{aligned}
S(q) &= \frac{L^2}{2\pi(l^*)^2 N} e^{-q^2 l^2/2} \\
& \sum_{\alpha, E_\alpha > 0} \langle \rho(\vec{q}) | \Psi_\alpha^R(\vec{q}) \rangle \langle \Psi_\alpha^L(\vec{q}) | \rho(\vec{q}) \rangle \quad (59)
\end{aligned}$$

where N is the number of particles, and we have defined

$$\langle \rho(\vec{q}) | \rangle_{m_1 m_2} = \rho_{m_1 m_2}(\vec{q}) \quad (60)$$

$$| \rho(\vec{q}) \rangle_{m_1 m_2} = (N_F(m_2) - N_F(m_1)) \rho_{m_2 m_1}(-\vec{q}) \quad (61)$$

Recalling the relation between the right and left eigenvectors eq(43), and noting that $N = pL^2/2\pi(l^*)^2$ for $\nu = \frac{p}{2p+1}$, we finally obtain

$$S(q) = \frac{1}{p} e^{-q^2 l^2/2} \sum_{\alpha, E_\alpha > 0} | \langle \rho(\vec{q}) | \Psi_\alpha^R(\vec{q}) \rangle |^2 \quad (62)$$

Our task in this section will be to compute all terms in this sum that contribute to order $(ql)^4$. It turns out that the case of a single CF-LL filled is nongeneric, while any $p > 1$ ($\nu = p/2p+1$) is generic. We address the simpler $p = 1$ case first.

A. The Structure factor for the Laughlin fractions

From the structure of eq(62) it is clear that only magnetoexciton operators with $N_F(m_1) \neq N_F(m_2)$ need to be kept in computing the density correlator. For $\nu = \frac{1}{3}$ this corresponds to O_{m0} (positive energy) and O_{0m} (negative energy). We will assume that the eigenvectors Ψ_α are analytic at small q , which turns out to be correct in the incompressible states. Furthermore, we will need the matrix elements ρ_{m0} which are

$$\rho_{m0}(\vec{q}) = \sqrt{\frac{1}{m!}} e^{im(\theta_q - \pi/2)} \left(\frac{cQ}{\sqrt{2}}\right)^m e^{-c^2 Q^2/4} \quad (63)$$

and $\rho_{0m}(\vec{q}) = (\rho_{m0}(-\vec{q}))^*$. It is clear that $\rho_{10} \approx q$, and $\rho_{20} \approx q^2$, and the other matrix elements go with higher powers of q . Therefore one needs to keep only the components 10, 20, 01, and 02 of all the eigenvectors Ψ_α to find $S(q)$ to order q^4 .

In order to compute these components of the eigenvectors, we need to analyse the small- q structure of $\mathcal{H}$ in greater detail. In the following, we will set $\theta_q = \pi/2$ for simplicity. This choice makes $\mathcal{H}$ real.

First consider the diagonal matrix elements.

$$\mathcal{H}(m0; m0; q) = \epsilon(m) - \epsilon(0) + \frac{v(q)}{2\pi(l^*)^2 \sqrt{m!n!}} e^{-Q^2/2} \left(\frac{c^2 Q^2}{2}\right)^m - \int \frac{sds}{(2\pi)} v(s) e^{-S^2/2} L_m\left(\frac{c^2 S^2}{2}\right) J_0(QS) \quad (64)$$

Let us recall that the differences of energies $\epsilon(m) - \epsilon(0)$ are given in terms of the potential (eq(21))

$$\begin{aligned} \epsilon(m) - \epsilon(0) &= \int \frac{d^2 s}{(2\pi)^2} v(s) e^{-s^2 l^2/2} (|\rho_{00}(\vec{s})|^2 - |\rho_{m0}(\vec{s})|^2) \\ &= \int \frac{sds}{(2\pi)} v(s) e^{-S^2/2} \left(1 - \frac{1}{m!} \left(\frac{c^2 S^2}{2}\right)^m\right) \end{aligned} \quad (65)$$

Consider $\mathcal{H}(10; 10, q)$. For small q the direct term contributes only in order q (for a long-range potential, $v(q \rightarrow 0) \simeq 1/q$), or order q^2 (for a short-range potential, $v(q \rightarrow 0) \simeq \text{const}$), while the energy difference and the exchange term contribute to order q^0 . Setting $J_0 \approx 1$ to get the order 1 contribution, we find that it *vanishes for an arbitrary potential!* The other diagonal matrix elements $m0$ for $m \neq 1$ show no such behavior. This means that the naive magnetoexciton O_{10} has zero energy as $q \rightarrow 0$. This signals us that it must be related to the constraint. The detailed manner in which it is related will be seen shortly.

As an aside we note that when one tries to construct CF-magnetoexciton wavefunctions in the standard manner by taking a IQH magnetoexciton, multiplying by the Laughlin-Jastrow factor, and projecting, one finds⁴⁵ that the IQH 10 magnetoexciton projects to zero as $q \rightarrow 0$. This seems to be directly related to what we find here, but since it is difficult to go back to electronic wavefunctions from the Hamiltonian theory, it is not clear how to extend this relationship further.

Let us now consider the off-diagonal elements of $\mathcal{H}$. There are two different types, those confined to the positive energy subspace, and those that go between the positive and negative energy subspaces.

$$\begin{aligned} \mathcal{H}(m0 : n0 : q) &= \frac{v(q)}{2\pi(l^*)^2 \sqrt{m!n!}} e^{-Q^2/2} \left(\frac{cQ}{\sqrt{2}}\right)^{m+n} \\ &- (-1)^{n < (m,n)+m} (\text{sgn}(n-m))^{n-m} \int \frac{sds}{(2\pi)} v(s) e^{-S^2/2} \\ &\times \sqrt{\frac{n_{<}(n,m)!}{n_{>}(n,m)!}} \left(\frac{cS}{\sqrt{2}}\right)^{|n-m|} L_{n_{<}}^{|n-m|} \left(\frac{c^2 S^2}{2}\right) J_{|n-m|}(QS) \end{aligned} \quad (66)$$

The most important property to note is that the off-diagonal matrix element vanishes as $q^{|n-m|}$ as $q \rightarrow 0$ (this behavior comes from the exchange term). Secondly, by exchanging m and n , it can be seen that $\mathcal{H}_{++}$ is a symmetric matrix. Now for the second kind of off-diagonal element

$$\begin{aligned} \mathcal{H}(m0; 0n; q) &= (-1)^n \frac{v(q)}{2\pi(l^*)^2 \sqrt{m!n!}} e^{-Q^2/2} \left(\frac{cQ}{\sqrt{2}}\right)^{m+n} \\ &- (-1)^n \int \frac{sds}{(2\pi)} v(s) e^{-S^2/2} \frac{J_{m+n}(QS)}{\sqrt{m!n!}} \end{aligned} \quad (67)$$

Once again, this vanishes as q^{m+n-1} (long-range potential) or q^{m+n} (short-range potential).

Thus, as $q \rightarrow 0$, the naive magnetoexcitons O_{m0} , O_{0m} become decoupled for $m > 1$, and therefore become the true eigenstates of $\mathcal{H}$. O_{10} and O_{01} have a diagonal energy of order q (long-range potential), and are coupled by matrix elements of order q , so they do not decouple. We will use these $q \rightarrow 0$ naive magnetoexcitons as labels for the true eigenstates, denoting the positive energy ones by Ψ_m and the negative energy ones by Ψ_{-m} (for $m > 1$). In order to find the true eigenstates Ψ_α at small q , one can do perturbation theory with q as the small parameter.

To see which of the true eigenstates we need to keep, consider the components of Ψ_3 . We have already seen that we need to restrict attention to the components 10, 20, 01, and 02, due to the powers of q already present in ρ . Due to the small q structure of $\mathcal{H}$ it can be seen that

$$\Psi_3^R \approx |30\rangle + \alpha q |20\rangle + \beta q |40\rangle + \gamma q^2 |10\rangle + \dots \quad (68)$$

where $|m0\rangle$ represent the naive magnetoexciton states. Therefore the overlap $\langle \rho(q) | \Psi_3^R \rangle \approx q^3$, and it is unnecessary to compute it if one wants $S(q)$ to order q^4 . Similar reasoning holds for Ψ_m for any $m > 2$ and $m < -2$.

To summarize, we see that to obtain $S(q)$ to order q^4 it is sufficient to compute $|\langle \rho(q) | \Psi_2 \rangle|^2$, since Ψ_1 will belong to the null (or unphysical) subspace. Also, the overlap needs to be computed only to an accuracy of q^2 .

To make further progress, let us write down the matrix $\mathcal{H}$ restricted to the lowest-energy four-dimensional subspace spanned by $|20\rangle$, $|10\rangle$, $|01\rangle$, and $|02\rangle$, where all the diagonal matrix elements have been kept to order q^2 , and a long-range interaction has been used.

$$\mathcal{H}_4 = \begin{pmatrix} E_{20}(k) & Bk & O(k^2) & O(k^3) \\ Bk & Ak & Ak - Ck^2 & O(k^2) \\ O(k^2) & -Ak + Ck^2 & -Ak & Bk \\ O(k^3) & O(k^2) & Bk & -E_{20}(k) \end{pmatrix} \quad (69)$$

We have defined $k = ql$. The various coefficients are (for the Coulomb interaction with the energy scale $e^2/\epsilon l$ removed, $v(q) = 2\pi l/q$)

$$\begin{aligned}
A &= \frac{1}{3} \\
B &= -\frac{1}{4}\sqrt{\frac{\pi}{3}} \\
C &= \frac{3}{8}\sqrt{\frac{\pi}{6}} \\
E_{20} &= \frac{1}{3}\sqrt{\frac{\pi}{6}} + O(q^2)
\end{aligned} \tag{70}$$

The central 2×2 submatrix consisting of the states 10 and 01 can be seen to be very singular. Taken by itself, it possesses two zero eigenvalues (up to order q^2) but only one eigenvector, and represents the constraint in its small- q form. The larger matrix also possesses two zero eigenvectors. This can be seen from the determinant of $\mathcal{H}_4$, which is

$$\det|\mathcal{H}_4| = 2AE_{20}k^3(CE_{20} - B^2) + k^4(B^4 - C^2E_{20}^2) \tag{71}$$

The key identity

$$B^2 = CE_{20}(0) \tag{72}$$

makes this determinant zero to order k^4 inclusive, which means that there are two zero eigenvalues to order k^2 . Eq(72) seems accidental, and dependent on the potential used, but it can be shown from the form of the low-order Laguerre polynomials that this identity holds independent of the potential, and furthermore that

$$B = -\frac{3}{2\sqrt{2}}E_{20}(0) \tag{73}$$

for any potential.

As mentioned before, the true eigenvectors can be found by straightforward perturbation theory. We first diagonalize the positive energy subspace by a unitary transformation (the Hamiltonian is hermitian in this subspace). The eigenvectors can be used to construct the unitary transformation which needs to be correct only to order k

$$U = \begin{pmatrix} 1 & -\frac{Bk}{E_{20}} \\ \frac{Bk}{E_{20}} & 1 \end{pmatrix} \tag{74}$$

The resulting eigenvalues are $E_{20} + \frac{B^2k^2}{E_{20}} \equiv E$ and $Ak - B^2k^2/E_{20} \equiv Ak - Ck^2$, and the matrix in the new basis looks like

$$U^\dagger \mathcal{H}_4 U = \begin{pmatrix} E & 0 & O(k^2) & O(k^3) \\ 0 & Ak - Ck^2 & Ak - Ck^2 & O(k^2) \\ O(k^2) & -Ak + Ck^2 & -Ak + Ck^2 & 0 \\ O(k^3) & O(k^2) & 0 & -E \end{pmatrix} \tag{75}$$

It is clear that to order q the nonzero positive eigenvector is decoupled from the negative energy subspace. The central 2×2 matrix is degenerate, and has two zero eigenvalues, but only one eigenvector. It represents the null subspace of $\mathcal{H}$ in the new basis. In order to compute $S(q)$ to order q^4 we need only the physical positive energy eigenvector Ψ_2 . It can be seen that Ψ_2 is decoupled from the null subspace. To the requisite order Ψ_2 is

$$\Psi_2^L = \begin{pmatrix} 1 & \frac{Bk}{E_{20}(0)} & O(k^2) & O(k^3) \end{pmatrix} \tag{76}$$

It can be easily verified that Ψ_2 is orthogonal to the null eigenvector, and this fact provides a shortcut to deriving the above results. Now taking the overlap with $|\rho\rangle$ and squaring, one easily finds

$$S(q) = \frac{(ql)^4}{8} + \dots \tag{77}$$

This coefficient differs by a factor of 2 from that predicted from Laughlin's wave function³ using the compressibility sum rule²², namely $S(q) = (ql)^4/4$. Therefore this calculation, while unambiguously showing the signature of an incompressible state, does not give the same result as the Laughlin state. This is a little surprising, since one expects the CF-HF state of one CF-LL filled to correspond to the Laughlin state. However, two points should be noted here. Firstly, it is very difficult to make the correspondence between the CF-wavefunction and the electronic one in the Hamiltonian formalism. It may be that the electronic wavefunction corresponding to one full CF-LL is incompressible, but not Laughlin's wavefunction. Secondly, this is an approximate calculation, albeit one that is conserving, and a more sophisticated conserving approximation may well produce the same answer as Laughlin's wave function.

Finally, one may extend this result to all the Laughlin fractions $\frac{1}{2s+1}$ by attaching $2s$ units of flux, with $c^2 = \sqrt{\frac{2s}{2s+1}}$ to obtain

$$S(q) = \frac{1}{8}(ql)^4 + \dots \tag{78}$$

independent of s , which also differs from the result for the coefficient obtained from Laughlin's wavefunction²², namely $s/4$.

It should be emphasized that the results are independent of the form of the potential. The short-range potential produces a slightly different form of the matrix $\mathcal{H}_4$ but one still obtains the same answer for the small q behavior of $S(q)$. One expects this since the structure factor is a property of the ground state, and should not depend on the dynamics.

B. The Structure Factor for the Principal Fractions $\nu = p/2p + 1$, $p > 1$

Let us now turn to the more generic case of $p > 1$. In this case we have p CF-LLs filled. We have already seen that only naive magnetoexcitons with $N_F(m_2) \neq N_F(m_1)$ contribute to the density correlator, so we will only consider those below. Once again, we find that the diagonal matrix element $\mathcal{H}(p, p-1; p, p-1; \vec{q})$, which is

$$\begin{aligned}
&\epsilon(p) - \epsilon(p-1) + \frac{v(q)}{2\pi(l^*)^{2p}} e^{-Q^2/2} \left(\frac{cQ}{\sqrt{2}}\right)^2 \\
&- \int \frac{sds}{2\pi} v(s) e^{-S^2/2} L_{p-1}\left(\frac{c^2 S^2}{2}\right) L_p\left(\frac{c^2 S^2}{2}\right) J_0(QS)
\end{aligned} \tag{79}$$

vanishes as $q \rightarrow 0$. Recall that the energy difference is (eq(21))

$$\epsilon(p) - \epsilon(p-1) = \int \frac{d^2 q}{(2\pi)^2} v(q) e^{-Q^2/2} \sum_{k=0}^{p-1} \left(\frac{k!}{(p-1)!} \left(\frac{c^2 Q^2}{2} \right)^{p-1-k} (L_k^{p-1-k})^2 - \frac{k!}{p!} \left(\frac{c^2 Q^2}{2} \right)^{p-k} (L_k^{p-k})^2 \right) \quad (80)$$

The $p, p-1$ diagonal matrix element vanishes as $q \rightarrow 0$ for an *arbitrary* potential by virtue of a Laguerre polynomial identity

$$\sum_{k=0}^{p-1} \frac{k!}{(p-1)!} x^{p-1-k} (L_k^{p-1-k}(x))^2 - \frac{k!}{p!} x^{p-k} (L_k^{p-k}(x))^2 = L_{p-1}(x) L_p(x) \quad (81)$$

This identity has been verified to high order p , but a general proof is not known to the author.

Once again, this vanishing energy is related to the constraint. Proceeding to the modes that are connected to this one by order q matrix elements, we find that there are two, $O_{p+1,p-1}$ and $O_{p,p-2}$. This is the primary difference between the case of generic p and $p = 1$, since in the latter case the second operator was missing. Now one proceeds as before, by first looking at the positive energy subspace

$$\mathcal{H}_{+,3} = \begin{pmatrix} E_1(q) & V_{12}(q) & B_1 k \\ V_{12}(q) & E_2(q) & B_2 k \\ B_1 k & b_2 k & E_0(q) \end{pmatrix} \quad (82)$$

where

$$E_0 = \mathcal{H}(p, p-1; p, p-1; q) \quad (83)$$

$$E_1 = \mathcal{H}(p+1, p-1; p+1, p-1; q) \quad (84)$$

$$E_2 = \mathcal{H}(p, p-2; p, p-2; q) \quad (85)$$

$$V_{12} = \mathcal{H}(p+1, p-1; p, p-2; q) \quad (86)$$

$$\mathcal{H}(p, p-1; p+1, p-1; q) = B_1 k + \dots \quad (87)$$

$$\mathcal{H}(p, p-1; p, p-2; q) = B_2 k + \dots \quad (88)$$

It should be noted that V_{12} does not vanish as $q \rightarrow 0$. Next one carries out a unitary transformation that makes the upper left 2×2 submatrix diagonal. This unitary transformation is

$$U_1 = \frac{1}{D} \begin{pmatrix} V_{12} & -(E_+ - E_1) & 0 \\ E_+ - E_1 & V_{12} & 0 \\ 0 & 0 & D \end{pmatrix} \quad (89)$$

where

$$E_{\pm} = \frac{1}{2}(E_1 + E_2 \pm \sqrt{(E_1 - E_2)^2 + V_{12}^2}) \quad (90)$$

$$D = \sqrt{(E_+ - E_1)^2 + V_{12}^2} \quad (91)$$

After this transformation the Hamiltonian in the 3×3 positive energy subspace assumes the form

$$\mathcal{H}_{+,3} = \begin{pmatrix} E_+ & 0 & k\tilde{B}_1 \\ 0 & E_- & k\tilde{B}_2 \\ k\tilde{B}_1 & k\tilde{B}_2 & 0 \end{pmatrix} \quad (92)$$

where

$$\tilde{B}_1 = \frac{V_{12}B_1 + (E_+ - E_1)B_2}{D} \quad (93)$$

$$\tilde{B}_2 = \frac{V_{12}B_2 - (E_+ - E_1)B_1}{D} \quad (94)$$

Let us call the basis vectors here Ψ_+ , Ψ_- , and Ψ_0 . Now it is easy to construct new basis vectors $\tilde{\Psi}_+$, $\tilde{\Psi}_-$ and $\tilde{\Psi}_0$ in the positive energy subspace that make it completely diagonal to order q^2 by doing first order perturbation theory in the off-diagonal elements. Once again it turns out that with this transformation, the positive energy subspace is decoupled from the negative energy and null subspaces to order q^2 . The structure factor is then computed from the overlaps of the density with the physical positive energy eigenvectors.

$$S(q) = \frac{1}{p} (|\langle \rho(q) | \tilde{\Psi}_+(q) \rangle|^2 + |\langle \rho(q) | \tilde{\Psi}_-(q) \rangle|^2) = (ql)^4 p \left[\left(\frac{1}{\sqrt{p+1}} + \frac{E_2 B_1 - V_{12} B_2}{E_1 E_2 - V_{12}^2} \right)^2 + \left(\frac{1}{\sqrt{p-1}} + \frac{E_1 B_2 - V_{12} B_1}{E_1 E_2 - V_{12}^2} \right)^2 \right] \quad (95)$$

All the above algebraic steps are heavily dependent on the potential used, but nevertheless one expects that the final result should be potential independent. Once again, a formal proof is lacking, but this potential-independence has been verified for the Coulomb potential and the short range Gaussian potential for $2/5$ and $3/7$. By some mysterious (to the author) properties of the Laguerre polynomials and their integrals, all the dependence on the functional form of the potential drops out.

Let us illustrate this with the Gaussian potential $v(q) = 2\pi(l^*)^2 e^{-\alpha Q^2/2}$, which is very convenient for explicitly carrying out integrals, but which (for large α) also possesses a small parameter $\beta = 1/(1 + \alpha)$ which will be exploited later. For $p = 2$ the relevant quantities (without any approximation) are

$$E_1 = \frac{12}{5}\beta^2 - \frac{32}{5}\beta^3 + \frac{192}{25}\beta^4 - \frac{2048}{625}\beta^5 \quad (96)$$

$$E_2 = \frac{4}{5}\beta^2 + \frac{32}{25}\beta^3 - \frac{192}{125}\beta^4 \quad (97)$$

$$V_{12} = -\frac{1}{\sqrt{3}} \left(\frac{12}{5}\beta^2 - \frac{96}{25}\beta^3 + \frac{192}{125}\beta^4 \right) \quad (98)$$

$$B_1 = -\frac{\beta^2}{\sqrt{3}} \left(3 - \frac{48}{5}\beta + \frac{336}{25}\beta^2 - \frac{768}{125}\beta^3 \right) \quad (99)$$

$$B_2 = \beta^2 \left(1 - \frac{16}{5}\beta + \frac{48}{25}\beta^2 \right) \quad (100)$$

The expression appearing in the denominator of eq(95) is

$$E_1 E_2 - V_{12}^2 = \beta^4 \left(\frac{2^9}{3^3}\beta - \frac{2^{13}}{3^4}\beta^2 + \frac{2^{16}}{3^5}\beta^3 - \frac{2^{18}}{3^6}\beta^4 + \frac{3 \times 2^{17}}{5^7}\beta^5 \right) \quad (101)$$

It can easily be verified that both $E_1 B_2 - V_{12} B_1$ and $E_2 B_1 - V_{12} B_2$ are proportional to the above expression, giving ratios independent of β . The structure factor depends on these ratios only, and is thus independent of β .

For $p > 3$ it becomes very difficult to compute E_i , B_i , and V_{12} exactly, even for the Gaussian potential. However, if one *assumes* the potential independence of $S(q)$, then one can use the following strategy: Since $S(q)$ does not depend on β , it can be calculated in the limit of small

β . In this limit, one need only compute terms of lowest nontrivial order in β in E_i , B_i , and V_{12} . Keeping one more order as a consistency check one finds for generic p

$$E_1 = \beta^2 c^2 (p+1) - 2\beta^3 c^4 (p^2 + p - 1) + \dots \quad (102)$$

$$E_2 = \beta^2 c^2 (p-1) - \beta^3 c^4 (2p^2 - 6p + 2) + \dots \quad (103)$$

$$V_{12} = -\sqrt{p^2 - 1}(\beta^2 c^2 - 2\beta^3 c^4 (p-1) + \dots) \quad (104)$$

$$B_1 = \sqrt{p+1}\beta^2(1 - \beta c^2(3p-2) + \dots) \quad (105)$$

$$B_2 = \sqrt{p-1}\beta^2(1 - \beta c^2(3p-2) \dots) \quad (106)$$

Using the above expressions in eq(95) one finds the result quoted in the introduction

$$S(q) = \frac{(ql)^4}{2} \frac{p^4 - 3p^3 + \frac{5}{4}p^2 + 3p + \frac{7}{4}}{p^2 - 1} \quad (107)$$

This expression, with its divergence as $p \rightarrow \infty$ or $\nu \rightarrow \frac{1}{2}$ is consistent with the result²⁶ that for a problem equivalent to the $\nu = \frac{1}{2}$ problem, $S(q) \simeq q^3 \log(q)$. As $p \rightarrow \infty$, the radius of convergence of the power series expansion of $S(q)$ must go to zero (else the structure factor would diverge for a range of q according to eq(107)), and the $p \rightarrow \infty$ limit does not commute with the $q \rightarrow 0$ limit.

Finally, it is interesting to compare the conserving prediction for the q^4 term of $S(q)$ with the naive calculation using the preferred density in the HF state

$$S_{naive}(q) = \langle \Omega_{HF} | \rho_p(\vec{q}) \rho_p(-\vec{q}) | \Omega_{HF} \rangle \quad (108)$$

This latter number turns out to be $p/8$. Read²⁶ wrote the conserving density-density correlator as a sum of the naive correlator of the preferred density and a term representing the coupling of this preferred density to a gauge field. Making a similar decomposition here we see that the gauge-field contribution increases with p , and comes to dominate the small- q structure factor as $p \rightarrow \infty$.

VI. EXPLICIT RESULTS FOR $\frac{1}{3}$ AND $\frac{2}{5}$

To motivate the results presented in this section let us consider the relation of the all- q Hamiltonian theory³⁵ to the original electronic theory. The all- q theory is algebraically consistent at all q and has the same algebra for the electronic density as the electronic theory. It is also consistent with the small- q theory that was developed earlier by canonically transforming the electronic Hamiltonian. If one could find a sequence of canonical transformations that exactly mapped the electronic Hamiltonian into this one, then one could be confident that the two problems are identical. In the absence of such a transformation, all we can really be sure of is that the all- q theory has the same small- q behavior as the electronic problem. It is therefore of interest to compute the magnetoexciton dispersions for q not very small to see if it agrees with the numerical results for the electronic problem. In this section we will present results for the spin-polarized magnetoexcitons for $\frac{1}{3}$ and $\frac{2}{5}$ and show that the minima in the lowest energy magnetoexciton dispersion occur at

the same values of ql as those found for the electronic problem.

As shown in the previous section, for very small q , the naive magnetoexcitons decouple and become the true eigenstates of $\mathcal{H}$. As q increases they become increasingly coupled, and both level repulsion (between positive energy states) and level attraction (between positive and negative energy states) manifest themselves, giving rise to complicated magnetoexciton dispersions. The matrix $\mathcal{H}$ is infinite-dimensional, but in any numerical calculation, only a finite matrix can be diagonalized. We saw that the lowest nontrivial result for $S(q)$ could be obtained by keeping at most a 6×6 matrix. As q increases more and more CF-LLs have to be kept to obtain accurate results. The accuracy of the results were checked by two different methods. Firstly, the number of CF-LLs kept was increased until the energy of the magnetoexciton was stable. Secondly, we know that at every q there should be two zero eigenvalues corresponding to the unphysical sector. For a finite number of CF-LLs kept, and q not too small, these would-be zero eigenvalues are not exactly zero, but much smaller than any other eigenvalues. The number of CF-LLs kept in the calculation was increased until these null eigenvalues were at least 4 orders of magnitude smaller than the smallest physical eigenvalue.




FIG. 1. Spin-polarized magnetoexciton dispersions for $\nu = \frac{1}{3}$ at $\lambda = 0.9l$ and $\lambda = l$. All energies are in units of $e^2/\epsilon l$.

The two figures are for the long-range potential $v(q) = \frac{2\pi e^2}{\epsilon q} e^{-q^2 \lambda^2/2}$ for various values of λ . It is worth noting that though the depths of the minima are λ -dependent, and different from what are found in numerical diagonalizations or from CF-wavefunctions, the positions are correct⁵⁻⁸. This indicates that the all- q theory does capture some of the important physics of the electronic problem even at fairly large q . Furthermore, the results are calculated in the lowest level conserving approximation. A more sophisticated conserving approximation incorporating the screening of the potential may produce better results at large q .




FIG. 2. Spin-polarized magnetoexciton dispersions for $\nu = \frac{2}{5}$ at $\lambda = 0$ and $\lambda = 0.5l$. All energies are in units of $e^2/\epsilon l$.

VII. EXTENSION OF THE FORMALISM TO NONUNIFORM DENSITY

In this section we will see how the fractional quasiparticle charge emerges from the formalism, provided flux attachment is carried out for the inhomogeneous problem. The all- q Hamiltonian theory³⁵ is based on the expectation that the ground state is uniform. In the presence of a localized quasiparticle this is clearly not a good assumption. More generally, it would be useful to have a conserving formulation which was applicable to the case of a nonuniform ground state, since this would enable a treatment of edges⁴⁶ or disorder.

To construct such a formalism, let us go back to the defining equations for the guiding center coordinates of the electron and the pseudovortex eq(6,7). Let us assume that the ground state density is $n(\vec{r}) = \bar{n} + \delta n(\vec{r})$. According to the principles of flux attachment, the CFs should see an additional effective gauge field corresponding to this nonuniform density which satisfies

$$\nabla \times \delta \vec{a} = 4\pi \delta n(\vec{r}) \quad (109)$$

This additional gauge field should be present in the CF-velocity operator $\vec{\Pi}$, leading to the commutation relation

$$[\Pi_{i\alpha}, \Pi_{i\beta}] = \frac{i\epsilon_{\alpha\beta}}{l^2} (1 - c^2 - 4\pi l^2 \delta n(\vec{r}_i)) \quad (110)$$

With this change, the quasiparticle charge can be computed by a very simple argument, identical in spirit to the counting argument given by Jain³⁸. Consider a region with large area A which contains one extra localized quasiparticle (in addition to p filled CF-LLs) in its interior. The total charge in this region (after subtracting the background) is

$$Q = -e\delta n(q \rightarrow 0) = -e(1 + \delta n_R(q \rightarrow 0)) \quad (111)$$

where the 1 counts the extra CF that has been introduced, and δn_R is the response of the filled CF-LLs to

the introduction of the extra CF. Before the extra CF was introduced, all the states in the CF-LLs $0, 1, \dots, p-1$ were occupied. If one assumes that the final state is adiabatically connected to the initial one, then the same states continue to be occupied. However, the total effective flux through the area A has decreased in magnitude by $4\pi\delta n(q \rightarrow 0)$. This means that each CF-LL has $2\delta n(q \rightarrow 0)$ fewer states in the area A than it did before. In essence, the states in the regions of smaller effective field have a larger cyclotron radius, so they expand and push out the other states. This means the total charge in the area A is

$$Q = -e\delta n(q \rightarrow 0) = -e(1 - 2p\delta n(q \rightarrow 0)) \quad (112)$$

which can be seen to give

$$Q = \frac{-e}{2p+1} \quad (113)$$

The inclusion of the varying gauge field in $\vec{\Pi}$ is essential for the success of this argument. Unfortunately, this modification of $\vec{\Pi}$ has an undesirable effect: Using the modified $\vec{\Pi}$ in the expressions for the guiding center coordinates $\vec{R}_e$ (eq(6)) and $\vec{R}_v$ (eq(7)) spoils their commutation relations (eqs(8,9,10)). This is a serious problem, since this means that the density operator constructed from the new $\vec{R}_e$ does not obey the magnetic translation algebra, and thus cannot be identified with the physical electronic density. The density also fails to commute with the constraint, making a conserving approximation infeasible.

However, there is a way to modify $\vec{R}_e$ and $\vec{R}_v$ so that the algebraic consistency of the theory is restored. Of the three commutation relations (eqs(8,9,10)) only the first two are essential to having a consistent theory. Satisfying eq(8) would mean that the density operator constructed from $\vec{R}_e$ could be identified with the electronic density, while satisfying eq(9) would mean that this density commutes with the constraint. If eq(10) is not satisfied it means that the constraint algebra does not close. New constraints would be generated by commuting old ones. This is not a problem for a conserving approximation, since by the Jacobi identity the electronic density operator (and thus the Hamiltonian constructed from it) would commute with the new constraints as well.

It turns out that operators $\vec{R}_e$, $\vec{R}_v$ which satisfy the conditions of the above paragraph to first order in $\frac{\delta n}{\bar{n}}$ can be constructed. They are

$$R_{e\alpha} = r_\alpha + \frac{l^2}{1+c} X_\alpha - \frac{2\pi l^4}{c(1+c)^2} \int_0^1 dt (1-t) \times (X_\alpha \delta n(\vec{r} - \frac{tl^2}{c(1+c)} \vec{X}) + \delta n(\vec{r} - \frac{tl^2}{c(1+c)} \vec{X}) X_\alpha) \quad (114)$$

$$R_{v\alpha} = r_\alpha - \frac{l^2}{c(1+c)} X_\alpha + \frac{2\pi l^4}{c^3(1+c)^2} \int_0^1 dt (c+t) \times (X_\alpha \delta n(\vec{r} - \frac{tl^2}{c(1+c)} \vec{X}) + \delta n(\vec{r} - \frac{tl^2}{c(1+c)} \vec{X}) X_\alpha) \quad (115)$$

where

$$X_\alpha = \epsilon_{\alpha\beta} \Pi_\beta \quad (116)$$

$$\delta n(\vec{r} - z\vec{X}) = \int \frac{d^2q}{(2\pi)^2} \delta n(\vec{q}) e^{i\vec{q} \cdot (\vec{r} - z\vec{X})} \quad (117)$$

and $\delta n(\vec{q})$ is the Fourier transform of the ground state expectation value of the electronic density operator, and is a commuting number. The $\vec{\Pi}$ operators appearing in the above formulas contain the spatially varying gauge field $\delta \vec{a}$. The preservation of the commutation relations eqs(8,9) to first order in $\delta n/n$ can be verified in a straightforward manner. This is therefore a good representation of the electronic problem for small deviations of density from a liquid state. Presumably eqs(114,115) can be extended to all orders in $\delta n/n$ while still maintaining the crucial commutation relations, eqs(8,9).

The modified density operator can in principle be used to calculate the detailed charge profile of a quasiparticle. One can create an extra CF in the p^{th} CF-LL, and solve the CFHF equations with the above expression for the density operator. There is a self-consistent aspect to the formulation: The $\delta n(\vec{q})$ that enters the density operator has to ultimately be the expectation value of the same density operator in the ground state.

The modifications to $\vec{R}_e$ and $\vec{R}_v$ to order $\delta n/n$ (other than the change in $\vec{\Pi}$) are not relevant for the charge of an isolated quasiparticle, but they are crucial if one wants to compute bosonic response functions for a nonuniform state in a conserving approximation. This issue will be of particular importance in the edge state problem⁴⁶. There one knows that there are *physical* gapless modes⁴⁶, and separating them from the unphysical modes can only be reliably carried out within a conserving approximation.

VIII. DISCUSSION AND CONCLUSIONS

The fractional quantum Hall states¹ have long had the peculiar status of being understood in principle^{3,4}, and yet being resistant to analytical calculation of their response functions. In the past few years much understanding has been achieved on how to compute these response functions. This understanding has largely been based on the notion of field-theoretic flux attachment via a Chern-Simons transformation¹⁰⁻¹⁶. Outstanding issues in the initial CS formulations¹⁷ were partially solved by Shankar and the author by adding variables and carrying out a sequence of canonical transformations to decouple the high and low energy degrees of freedom^{18,19}. However, the resulting theory, though it had many desirable features at small q , was unsatisfactory when viewed as a theory of the Lowest Landau level. Shankar posited a formulation³⁵ which was algebraically consistent at all q as a LLL theory, and had all the desirable features of the small- q formulation.

In the past, analysis of this all- q formalism³⁵ was limited to the HF approximation in which one used the preferred density $\rho_p = \rho_e - c^2 \rho_v$. The preferred density makes explicit many nonperturbative properties of the true density operator, such as the fractional quasiparticle charge and the order q^2 matrix elements out of the

ground state for incompressible states. The price paid is that the constraints, once they are used to construct ρ_p , are ignored, and one is not sure how robust the results are under a proper implementation of the constraints.

The most important objective achieved in this paper is that the nonperturbative features that were introduced by hand into the preferred density¹⁹ are seen to emerge very naturally from the theory when constraints are consistently imposed and flux attachment is carried out for states of nonuniform density.

The conserving approximation used is identical in spirit to the one used by Read²⁶ to obtain response functions for the $\nu = 1$ boson problem based on the formalism of Pasquier and Haldane²⁵. The case considered here is in some ways more complicated in that it involves Landau level states instead of plane wave states, but also in other ways simpler, in that finite-dimensional matrix equations have to be solved to obtain the response rather than integral equations²⁶.

The order q^2 matrix elements of the charge density operator emerge when unphysical states are discarded and only physical states are considered. The crux of the matter is that for $\nu = \frac{p}{2p+1}$ only the $p-1$ to p transition can produce an order q matrix element in violation of Kohn's theorem, and as $q \rightarrow 0$ this magnetoexciton state becomes identical to the constraint to order q , and thus unphysical.

Once the formalism is extended to include variations of density in the ground state, the fractional quasiparticle charge^{3,37,38} also emerges from the theory.

Finally, a look at the spin-polarized magnetoexciton dispersions for q not too small shows that though one does not yet have a rigorous connection between the all- q theory and the original electronic theory, the all- q conserving approximation manages to capture much of the important physics, even at large q .

In summary, it appears that the all- q Hamiltonian formulation of the fractional quantum Hall problem coupled to a conserving approximation is well-suited to computing bosonic response functions in the spin-polarized case. An analysis of the edge and disordered problems also seems possible in this approach using the extension to small density variations presented in Section VII, and will be pursued in future work.

IX. ACKNOWLEDGEMENTS

I would like to thank J.K.Jain and R.Shankar for many helpful conversations, and the NSF for partial support (DMR-0071611).

X. APPENDIX I

Most of the material in this appendix appears in ref.[³¹], but is presented here to make the discussion self-contained. In order to compute the sums over CF-LL indices of products of ρ_e or ρ_v matrix elements, it turns

out to be convenient to express all the coordinates in terms of the guiding center and cyclotron coordinates of the CF, which are

$$\vec{R}_{CF}^g = \vec{r} - \frac{l^2}{(1-c^2)} \hat{z} \times \vec{\Pi} \quad (118)$$

$$\vec{R}_{CF}^c = \frac{l^2}{(1-c^2)} \hat{z} \times \vec{\Pi} \quad (119)$$

which have the commutation relations

$$[R_{CF,\alpha}^c, R_{CF,\beta}^c] = i \frac{l^2}{(1-c^2)} \epsilon_{\alpha\beta} \quad (120)$$

$$[R_{CF,\alpha}^g, R_{CF,\beta}^g] = -i \frac{l^2}{(1-c^2)} \epsilon_{\alpha\beta} \quad (121)$$

$$[R_{CF,\alpha}^c, R_{CF,\beta}^g] = 0 \quad (122)$$

We can now express the electron and pseudovortex guiding center coordinates in terms of the CF operators $\vec{R}_{CF}^c$ and $\vec{R}_{CF}^g$ as

$$\vec{R}_e = \vec{R}_{CF}^g + c\vec{R}_{CF}^c \quad (123)$$

$$\vec{R}_v = \vec{R}_{CF}^g + \frac{1}{c}\vec{R}_{CF}^c \quad (124)$$

The CF guiding center and cyclotron coordinates, in turn, can be expressed in terms of two sets of canonical, commuting creation and annihilation operators

$$R_{CF,x}^c + iR_{CF,y}^c = l\sqrt{\frac{2}{1-c^2}} a_c \quad (125)$$

$$R_{CF,x}^c - iR_{CF,y}^c = l\sqrt{\frac{2}{1-c^2}} a_c^\dagger \quad (126)$$

$$R_{CF,x}^g + iR_{CF,y}^g = l\sqrt{\frac{2}{1-c^2}} a_g^\dagger \quad (127)$$

$$R_{CF,x}^g - iR_{CF,y}^g = l\sqrt{\frac{2}{1-c^2}} a_g \quad (128)$$

where

$$[a_c, a_c^\dagger] = [a_g, a_g^\dagger] = 1 \quad (129)$$

$$[a_c, a_g^\dagger] = [a_g, a_c^\dagger] = 0 \quad (130)$$

We can abstractly set up the states of the CF-cyclotron creation and annihilation operators

$$|n\rangle = \frac{(a_c^\dagger)^n}{\sqrt{n!}} |0\rangle \quad (131)$$

In terms of these states the matrix elements of eq(19) and eq(20) are simply expressed as

$$\rho_{m_1 m_2}(\vec{q}) = \langle m_1 | e^{-ic\vec{q} \cdot \vec{R}_{CF}^c} | m_2 \rangle \quad (132)$$

$$\chi_{m_1 m_2}(\vec{q}) = \langle m_1 | e^{-\frac{i}{c}\vec{q} \cdot \vec{R}_{CF}^c} | m_2 \rangle \quad (133)$$

In other words, only the cyclotron part of the electron and pseudovortex density enters the matrix elements defined above. An important result we will need is the following sum

$$\sum_m \chi_{m_1 m}(\vec{q}_1) \rho_{m m_2}(\vec{q}_2) \quad (134)$$

Using the completeness of the cyclotron states $|m\rangle$ we can recast this sum as

$$\langle m_1 | e^{-\frac{i}{c}\vec{q}_1 \cdot \vec{R}_{CF}^c} e^{-ic\vec{q}_2 \cdot \vec{R}_{CF}^c} | m_2 \rangle \quad (135)$$

We can then use the commutation relations of eq(120) to combine the exponentials and obtain

$$\sum_m \chi_{m_1 m}(\vec{q}_1) \rho_{m m_2}(\vec{q}_2) = e^{-\frac{i}{2}\vec{q}_1 \times \vec{q}_2} \times \langle m_1 | e^{-i(\frac{\vec{q}_1}{c} + c\vec{q}_2) \cdot \vec{R}_{CF}^c} | m_2 \rangle \quad (136)$$

Finally we separate the exponentials in the reverse order to get the second useful identity

$$\begin{aligned} & \sum_m \chi_{m_1 m}(\vec{q}_1) \rho_{m m_2}(\vec{q}_2) \\ &= e^{-i\vec{q}_1 \times \vec{q}_2} \sum_m \rho_{m_1 m}(\vec{q}_2) \chi_{m m_2}(\vec{q}_1) \end{aligned} \quad (137)$$

Note that the cyclotron parts of ρ_e and ρ_v *do not commute*, even though the entire operators, including the CF-guiding center parts, do.

-
- ¹ D.Tsui, H.Stromer and A.Gossard, Phys. Rev. Lett. **48**, 1599, (1982).
 - ² For a broad survey of the quantum Hall effects, see, “*Perspectives in Quantum Hall Effects*”, S.Das Sarma and A.Pinczuk, Editors (Wiley Interscience, New York 1997).
 - ³ R.B.Laughlin, Phys. Rev. Lett. **50**, 1395, (1983)
 - ⁴ J.K.Jain, Phys. Rev. Lett. **63**, 199, (1989); Phys. Rev. **B41**, 7653 (1990); Science **266**, 1199 (1994).
 - ⁵ J.K.Jain and R.K.Kamilla, in Chapter 1 of, “*Composite Fermions*”, Olle Heinonen, Editor (World Scientific, Teaneck, NJ, 1998).
 - ⁶ F.D.M.Haldane and E.H.Rezayi, Phys. Rev. Lett. **54**, 237 (1985).
 - ⁷ N.d’Ambrumenil and R.Morf, Phys. Rev. **B40**, 6108 (1989); P.M.Platzman and S.He, Phys. Rev. **B49**, 13674 (1994); S.He, S.H.Simon, and B.I.Halperin, Phys. Rev. **B50**, 1823 (1994).
 - ⁸ X.G.Wu and J.K.Jain, Phys. Rev. **B51**, 1752 (1995); R.K.Kamilla, X.G.Wu, and J.K.Jain, Phys. Rev. Lett. **76**, 1332 (1996); Phys. Rev. **B54**, 4873 (1996).
 - ⁹ J.M.Leinaas and J.Myrheim, Nuovo Cimento **37B**, 1 (1977).
 - ¹⁰ S.M.Girvin, in Chapter 9 of, “*The Quantum Hall Effect*”, Edited by R.E.Prange and S.M. Girvin, Springer-Verlag, 1990.
 - ¹¹ S.M.Girvin and A.H.MacDonald, Phys. Rev. Lett. **58**, 1252 (1987).
 - ¹² S.-C.Zhang, H.Hansson and S.A.Kivelson, Phys. Rev. Lett. **62**, 82, (1989); D.-H.Lee and S.-C.Zhang, Phys. Rev. Lett. **66**, 1220 (1991); S.-C.Zhang, Int. J. Mod. Phys., **B6**, 25 (1992).
 - ¹³ N.Read, Phys. Rev. Lett., **62**, 86 (1989).
 - ¹⁴ A.Lopez and E.Fradkin, Phys. Rev. B **44**, 5246 (1991), *ibid* **47**, 7080, (1993), Phys. Rev. Lett. **69**, 2126 (1992).
 - ¹⁵ V.Kalmeyer and S.-C.Zhang, Phys. Rev. B **46**, 9889 (1992).

- ¹⁶ B.I.Halperin, P.A.Lee and N.Read, Phys. Rev. B **47**, 7312 (1993).
- ¹⁷ S.H.Simon, A.Stern, and B.I.Halperin, Phys. Rev. B **54**, R11114 (1996); S.H.Simon and B.I.Halperin, Phys. Rev. B **48**, 17368, (1993).
- ¹⁸ R.Shankar and G.Murthy, Phys. Rev. Lett. **79**, 4437, (1997).
- ¹⁹ G.Murthy and R.Shankar, in “*Composite Fermions*”, Olle Heinonen, Editor, (World Scientific, Teaneck, NJ, 1998).
- ²⁰ D.Bohm and D.Pines, Phys. Rev. **92**, 609, (1953).
- ²¹ W.Kohn, Phys. Rev **123**, 1242 (1961).
- ²² S.M.Girvin, A.H.MacDonald and P.Platzman, Phys. Rev. B **33**, 2481, (1986).
- ²³ N.Read, Semi. Sci. Tech. **9**, 1859 (1994); Surf. Sci., **361/362**, 7 (1996).
- ²⁴ D.H.Lee, Phys. Rev. Lett.**80**, 4745 (1998).
- ²⁵ V.Pasquier and F.D.M.Haldane, Nucl. Phys. **B516**, 719 (1998).
- ²⁶ N.Read, Phys. Rev. **B58**, 16262 (1998).
- ²⁷ A.Stern, B.I.Halperin, F.von Oppen, and S.H.Simon, Phys. Rev. **B 59**, 12547 (1999); F. von Oppen, B. I. Halperin, S. H. Simon, and A. Stern, cond-mat/9903430.
- ²⁸ G.Murthy and R.Shankar, Phys. Rev. **B 59**, 12260 (1999).
- ²⁹ G.Murthy, K.Park, R.Shankar, and J.K.Jain, Phys. Rev. **B 58**, 15363 (1998).
- ³⁰ G. Murthy, Jour. Phys. Cond. Mat. **12**, 10543 (2000).
- ³¹ R. Shankar, Phys. Rev. **B 63** 085322 (2001).
- ³² G. Murthy, Phys. Rev. **B 60**, 13702 (1999).
- ³³ F.F.Fang and W.E.Howard, Phys. Rev. Lett. **16**, 797 (1966); T.Ando, A.B.Fowler, and F.Stern, Rev. Mod. Phys. **54**, 437 (1982).
- ³⁴ F.C.Zhang and S.Das Sarma, Phys. Rev. B **33**, 2903 (1986); D.Yoshioka, J. Phys. Soc. Jpn. **55**, 885 (1986); T.Chakraborty, P.Pietlainen, and F.C.Zhang, Phys. Rev. Lett, **57**, 130 (1986); R.Morf and B.I.Halperin, Z. Phys. B **68**, 391 (1987); R.Price and S.Das Sarma, Phys. Rev. B **54**, 8033 (1996).
- ³⁵ R. Shankar, Phys. Rev. Lett. **83** 2382 (1999).
- ³⁶ G.Baym and L.P.Kadanoff, Phys. Rev. **124**, 287 (1961); L. P. Kadanoff and G. Baym, “*Quantum Statistical Mechanics*”, Addison-Wesley, Reading, MA, 1989.
- ³⁷ F.D.M.Haldane, Phys. Rev. Lett. **51**, 605 (1983); B.I.Halperin, Phys. Rev. Lett. **52**, 1583 (1984); W.P.Su, Phys. Rev. **B 30**, 1069 (1984).
- ³⁸ J.K.Jain, Comm. Cond. Matt. Phys. **16**, 307 (1993); A.S.Goldhaber and J.K.Jain, Phys. Lett. **A199**, 267 (1995).
- ³⁹ B.I.Halperin and A.Stern, Phys. Rev. Lett.**80**, 5457 (1998); G.Murthy and R.Shankar, *ibid* 5458 (1998).
- ⁴⁰ K.W.Chiu and J.J.Quinn, Phys. Rev. B **9**, 4724 (1974); N.Horing and M.Yildiz, Ann. Phys. NY **97**, 216 (1976); T.Theis, Surf. Sci. **98**, 515 (1980); P.Hawrylak and J.J.Quinn, Phys. Rev. **B31**, 6592 (1985); Yu.A.Bychkov, S.V.Iordanskii, and G.M.Eliashberg, JETP Lett. **33**, 152 (1981).
- ⁴¹ C.Kallin and B.I.Halperin Phys. Rev B **30**, 5655, (1984).
- ⁴² A.H.MacDonald, Jour. Phys. **C18**, 1003 (1985).
- ⁴³ H.C.A.Oji and A.H.MacDonald, Phys. Rev. **B33**, 3810 (1986); I.K.Marmorkos and S.Das Sarma, Phys. Rev. **B45**, 13396 (1992); J.P.Longo and C.Kallin, Phys. Rev. **B47**, 4429 (1993).
- ⁴⁴ G.Rickayzen, Phys. Rev. **115**, 795 (1957).
- ⁴⁵ J.K.Jain, private communication.
- ⁴⁶ For a general introduction to edge state theory see, C.L.Kane and M.P.A.Fisher in, “*Perspectives in Quantum Hall Effects*”, S.Das Sarma and A.Pinczuk, Editors (Wiley Interscience, New York 1997).